

COPY

4

ETHEL PERCY ANDRUS GERONTOLOGY CENTER
(213) 743 6060



AD-A213 100

September 27, 1989

Dr. Jeannine A. Majde
Code 1141SB
Office of Naval Research
800 N. Quincy Street
Arlington, VA 22217

RE: N00014-89-J-1005 Annual Report

Dear Dr. Majde:

Enclosed are three copies of the Annual Progress Report for "Steroids, Stress, and Gene Expression in Neurons". If you need additional Information please contact me at (213) 743-5168.

Sincerely,

Caleb E. Finch, Ph.D.
Principal Investigator

and

Nancy R. Nichols, Ph.D.
Co-Principal Investigator

Connie Whitley
Sr. Contract and Grant
Administrator

DTIC
ELECTE
OCT 03 1989
S DCS D

DISTRIBUTION STATEMENT A

Approved for public release;
Distribution Unlimited

cc: Distribution List

89 10 2 120

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

1a. REPORT SECURITY CLASSIFICATION			1b. RESTRICTIVE MARKINGS		
2a. SECURITY CLASSIFICATION AUTHORITY			3. DISTRIBUTION/AVAILABILITY OF REPORT		
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE			Distribution Unlimited		
4. PERFORMING ORGANIZATION REPORT NUMBER(S) University of Southern California			5. MONITORING ORGANIZATION REPORT NUMBER(S)		
6a. NAME OF PERFORMING ORGANIZATION University of Southern Calif.	6b. OFFICE SYMBOL (If applicable)	7a. NAME OF MONITORING ORGANIZATION Office of Naval Research			
6c. ADDRESS (City, State, and ZIP Code) University of Southern California Ethel Percy Andrus Gerontology Center, MC0191 Los Angeles, CA 90089-0191		7b. ADDRESS (City, State, and ZIP Code) 800 N. Quincy Street Arlington, VA 22217-5000			
8a. NAME OF FUNDING/SPONSORING ORGANIZATION Office of Naval Research	8b. OFFICE SYMBOL (If applicable) ONR	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER N00014-89-J-1005			
8c. ADDRESS (City, State, and ZIP Code) 800 N. Quincy Street Arlington, VA 22217-5000		10. SOURCE OF FUNDING NUMBERS			
		PROGRAM ELEMENT NO. 61153N	PROJECT NO. RR04108	TASK NO.	WORK UNIT ACCESSION NO.
11. TITLE (Include Security Classification) Steroids, Stress, and Gene Expression in Neurons					
12. PERSONAL AUTHOR(S) Dr. Caleb E. Finch and Dr. Nancy R. Nicholas					
13a. TYPE OF REPORT ANNUAL	13b. TIME COVERED FROM 10/88 TO 9/89	14. DATE OF REPORT (Year, Month, Day) 89/10/01		15. PAGE COUNT 2	
16. SUPPLEMENTARY NOTATION					
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)		
FIELD	GROUP	SUB-GROUP			
19. ABSTRACT (Continue on reverse if necessary and identify by block number)					
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED/UNLIMITED <input type="checkbox"/> SAME AS RPT. <input type="checkbox"/> DTIC USERS			21. ABSTRACT SECURITY CLASSIFICATION		
22a. NAME OF RESPONSIBLE INDIVIDUAL Dr. J. A. Maide		22b. TELEPHONE (Include Area Code) 202 696-6055		22c. OFFICE SYMBOL ONR	

DISTRIBUTION LIST

Dr. Jeannine A. Majde (3)
Code 1141SB
Office of Naval Research
800 N. Quincy Street
Arlington, VA 22217

Mr. Robert Bachman (1)
University of California, San Diego
MC A-034
La Jolla, CA 92093-0234

Director Naval Research Laboratory (6)
Attention 2627
Washington D.C. 20375

Defense Technical Information Center (12)
Building 5
Cameron Street
Arlington, VA 22314



1. Reason for _____
 2. YES _____ ✓
 3. NO _____
 4. _____
 5. _____
 6. _____
 7. _____
 8. _____
 9. _____
 10. _____
 11. _____
 12. _____
 13. _____
 14. _____
 15. _____
 16. _____
 17. _____
 18. _____
 19. _____
 20. _____
 21. _____
 22. _____
 23. _____
 24. _____
 25. _____
 26. _____
 27. _____
 28. _____
 29. _____
 30. _____
 31. _____
 32. _____
 33. _____
 34. _____
 35. _____
 36. _____
 37. _____
 38. _____
 39. _____
 40. _____
 41. _____
 42. _____
 43. _____
 44. _____
 45. _____
 46. _____
 47. _____
 48. _____
 49. _____
 50. _____
 51. _____
 52. _____
 53. _____
 54. _____
 55. _____
 56. _____
 57. _____
 58. _____
 59. _____
 60. _____
 61. _____
 62. _____
 63. _____
 64. _____
 65. _____
 66. _____
 67. _____
 68. _____
 69. _____
 70. _____
 71. _____
 72. _____
 73. _____
 74. _____
 75. _____
 76. _____
 77. _____
 78. _____
 79. _____
 80. _____
 81. _____
 82. _____
 83. _____
 84. _____
 85. _____
 86. _____
 87. _____
 88. _____
 89. _____
 90. _____
 91. _____
 92. _____
 93. _____
 94. _____
 95. _____
 96. _____
 97. _____
 98. _____
 99. _____
 100. _____
 101. _____
 102. _____
 103. _____
 104. _____
 105. _____
 106. _____
 107. _____
 108. _____
 109. _____
 110. _____
 111. _____
 112. _____
 113. _____
 114. _____
 115. _____
 116. _____
 117. _____
 118. _____
 119. _____
 120. _____
 121. _____
 122. _____
 123. _____
 124. _____
 125. _____
 126. _____
 127. _____
 128. _____
 129. _____
 130. _____
 131. _____
 132. _____
 133. _____
 134. _____
 135. _____
 136. _____
 137. _____
 138. _____
 139. _____
 140. _____
 141. _____
 142. _____
 143. _____
 144. _____
 145. _____
 146. _____
 147. _____
 148. _____
 149. _____
 150. _____
 151. _____
 152. _____
 153. _____
 154. _____
 155. _____
 156. _____
 157. _____
 158. _____
 159. _____
 160. _____
 161. _____
 162. _____
 163. _____
 164. _____
 165. _____
 166. _____
 167. _____
 168. _____
 169. _____
 170. _____
 171. _____
 172. _____
 173. _____
 174. _____
 175. _____
 176. _____
 177. _____
 178. _____
 179. _____
 180. _____
 181. _____
 182. _____
 183. _____
 184. _____
 185. _____
 186. _____
 187. _____
 188. _____
 189. _____
 190. _____
 191. _____
 192. _____
 193. _____
 194. _____
 195. _____
 196. _____
 197. _____
 198. _____
 199. _____
 200. _____
 201. _____
 202. _____
 203. _____
 204. _____
 205. _____
 206. _____
 207. _____
 208. _____
 209. _____
 210. _____
 211. _____
 212. _____
 213. _____
 214. _____
 215. _____
 216. _____
 217. _____
 218. _____
 219. _____
 220. _____
 221. _____
 222. _____
 223. _____
 224. _____
 225. _____
 226. _____
 227. _____
 228. _____
 229. _____
 230. _____
 231. _____
 232. _____
 233. _____
 234. _____
 235. _____
 236. _____
 237. _____
 238. _____
 239. _____
 240. _____
 241. _____
 242. _____
 243. _____
 244. _____
 245. _____
 246. _____
 247. _____
 248. _____
 249. _____
 250. _____
 251. _____
 252. _____
 253. _____
 254. _____
 255. _____
 256. _____
 257. _____
 258. _____
 259. _____
 260. _____
 261. _____
 262. _____
 263. _____
 264. _____
 265. _____
 266. _____
 267. _____
 268. _____
 269. _____
 270. _____
 271. _____
 272. _____
 273. _____
 274. _____
 275. _____
 276. _____
 277. _____
 278. _____
 279. _____
 280. _____
 281. _____
 282. _____
 283. _____
 284. _____
 285. _____
 286. _____
 287. _____
 288. _____
 289. _____
 290. _____
 291. _____
 292. _____
 293. _____
 294. _____
 295. _____
 296. _____
 297. _____
 298. _____
 299. _____
 300. _____
 301. _____
 302. _____
 303. _____
 304. _____
 305. _____
 306. _____
 307. _____
 308. _____
 309. _____
 310. _____
 311. _____
 312. _____
 313. _____
 314. _____
 315. _____
 316. _____
 317. _____
 318. _____
 319. _____
 320. _____
 321. _____
 322. _____
 323. _____
 324. _____
 325. _____
 326. _____
 327. _____
 328. _____
 329. _____
 330. _____
 331. _____
 332. _____
 333. _____
 334. _____
 335. _____
 336. _____
 337. _____
 338. _____
 339. _____
 340. _____
 341. _____
 342. _____
 343. _____
 344. _____
 345. _____
 346. _____
 347. _____
 348. _____
 349. _____
 350. _____
 351. _____
 352. _____
 353. _____
 354. _____
 355. _____
 356. _____
 357. _____
 358. _____
 359. _____
 360. _____
 361. _____
 362. _____
 363. _____
 364. _____
 365. _____
 366. _____
 367. _____
 368. _____
 369. _____
 370. _____
 371. _____
 372. _____
 373. _____
 374. _____
 375. _____
 376. _____
 377. _____
 378. _____
 379. _____
 380. _____
 381. _____
 382

ABSTRACT

In order to identify glucocorticoid responses that mediate cellular adaptation to stress, we have examined changes in gene expression in rat hippocampus in response to acute corticosterone (CORT) treatment or to vibratory stress. Previously, we described select changes in four poly(A)-containing RNAs, coding for 50, 35, 33 and 20 kd polypeptides, in response to administered CORT. We have cloned five CORT-responsive mRNAs from a rat hippocampal cDNA library by differential hybridization and compared them to the previous responses by hybrid selection, in vitro translation, and 2-d gel analysis. Two of the clones hybridized to mRNAs coding for the 50 and 35 kd polypeptides; they were identified from partial DNA sequences as glial fibrillary acidic protein and glycerol phosphate dehydrogenase. Another clone was identified as glutamine synthetase and two were not identified, CR16 and CR62. CORT and stress responsiveness of these mRNAs were compared by 2-d gel analysis of in vitro translation products or RNA hybridization experiments. Only GPDH/35 kd and 20 kd were responsive to 2 hr CORT treatment and 2 hr vibratory stress. The lack of response to stress of the other mRNAs, despite elevated CORT levels, is consistent with their slower response to exogenous CORT. One exception is 33 kd which does respond to 2 hr CORT treatment but not to 2 hr vibratory stress. Therefore, we find subsets of mRNA responses which differ in temporal and other aspects of CORT regulation. Examining a variety of stressors and chronic stress will determine if this is a general or unique pattern of expression of CORT-dependent responses in hippocampus.

INTRODUCTION

Glucocorticoids mediate cellular adaptation to stress by a variety of mechanisms including receptor-mediated changes in gene expression (Munck et al., 1984). Most studies in brain have focussed on neuropeptides or peptide hormones that regulate the hypothalamic-pituitary-adrenal (HPA) axis and hence the output of glucocorticoids (Swanson et al., 1986; Dallman et al., 1987). Additional mechanisms for adaptation to stress might be revealed by examining patterns of gene expression in brain in response to elevated circulating glucocorticoid levels. Since the hippocampus is both a glucocorticoid target tissue with a high concentration of receptors and a site of negative feedback regulation of the hypothalamic-pituitary-adrenal axis (McEwen et al., 1986; Ratka et al., 1989), we have examined changes in gene expression in rat hippocampus in response to corticosterone (CORT) and physiological stress.

Previously, we described select changes in four poly(A)-containing RNAs, coding for 50, 35, 33 and 20 kd polypeptides, in response to administered CORT (Nichols et al., 1988a). In addition, we have cloned five CORT-responsive mRNAs from a rat hippocampal cDNA library by differential hybridization (Masters et al., 1987; Nichols et al., in press). In this report we characterize (and in some cases, identify) the cloned mRNAs and compare them to the previous mRNA translation products. In addition, we have analyzed these CORT-responses for responsiveness to 2 hr vibratory stress. Based on these data, we discuss how these molecular markers may be used to investigate mechanisms of cellular adaptation to stress.

METHODS

Animals, Treatments, RNA Isolation and CORT RIA

Male Fischer 344 rats (200-250 g) were multiply housed with a controlled light-dark cycle (lights on 06.00 h and off 18.00 h). Food and water were ad libitum. Rats were either adrenalectomized (ADX) and maintained on 0.9% saline or intact; ADX rats were injected subcutaneously with CORT or Mazola corn oil vehicle. Following decapitation (2 or 8 h after a morning injection or 2 h after initiation of stress), hippocampi were rapidly dissected and stored at -80°C until total RNA was isolated from pooled or individual samples by a guanidinium thiocyanate/CsCl procedure (Nichols et al., 1988a). Trunk blood was collected for measurement of serum CORT by radioimmunoassay (Nichols et al., 1989).

Isolation and Identification of Rat Hippocampal cDNA Clones

A rat hippocampal cDNA library was constructed in λ gt10 from 1 μ g of poly(A)-containing RNA from both ADX and ADX + CORT (10 mg/day for 3 days) groups. Approximately 100,000 plaques were screened on replicate nitrocellulose filters by differential hybridization using 32 P-cDNA prepared from 0.5 μ g of each poly(A)-containing RNA pool used to make the library. CORT-responsive cDNA clones were rescreened twice, subcloned into the Bluescribe or Bluescript vectors (Stratagene) and partial DNA sequences were compared with known genes in GenBank (Nichols et al., in press).

Hybrid Selection, In Vitro Translation and Two-dimensional Gel Analysis

Rat brain mRNAs were selected from total RNA using immobilized cDNA clones and the hybrid selection method of Mason and Williams (1985). Total RNA or select RNA was used as a template for in vitro translation by a rabbit reticulocyte system with 35 S-methionine (Nichols et al.,

in press). Translation products were resolved by two-dimensional (2-d) gel electrophoresis, visualized by fluorography and quantitated by computerized videodensitometry (Nichols et al., 1988a).

RNA Hybridization Analysis

Total RNA from treated and control individual or pooled samples were either fractionated on formaldehyde (0.2M)/agarose gels and transferred to positively charged nylon membranes (Nichols et al., in press) or vacuum-blotted directly on nylon membranes using a slot-blot apparatus. Anti-sense ^{32}P -cRNA probes were transcribed from linearized subclones in a plasmid vector and random-primed ^{32}P -cDNA probes were made from gel-purified inserts. Hybridization procedures and quantitation of autoradiographs were as described in Nichols et al. (in press). Sizes of transcripts were determined by comparison with an RNA ladder stained with ethidium bromide. For more precise quantitation, RNA was titrated by the solution hybridization/RNase protection method of Lee and Costlow (1987) and expressed as pg transcript/ μg total RNA.

RESULTS

We have used two methods to examine altered patterns of gene expression in rat hippocampus in response to administered glucocorticoids: analysis of RNA in vitro translation products on 2-d gels (Nichols et al., 1988a) and cloning of cDNAs from a rat hippocampal library by differential hybridization (Nichols et al., in press). We have further characterized CORT-responsive cDNA clones in hybrid selection/in vitro translation experiments and compared physico-chemical characteristics of the polypeptide products (and in two cases, co-migration on 2-d gel fluorographs) with previously characterized CORT-responsive mRNA translation products (Table 1). From these data and DNA sequence analysis (Table 1), two of the four mRNA translation products have been identified as glycerol phosphate dehydrogenase (GPDH)/35 kd and glial fibrillary acidic protein (GFAP)/50 kd; two others are still unidentified, 33 kd and 20 kd. An additional clone was identified as glutamine synthetase (GS/43 kd) and two were not identified, CR16 (not resolved on 2-d gels) and CR62/43 kd.

Based on their rapid response (2 h) of large magnitude (5- to 50-fold) to high doses of CORT (50 mg/kg/day) and type II glucocorticoid receptor-specificity, we postulated that the hippocampal mRNA increases should also occur during response to stress-induced elevations of glucocorticoid (Nichols et al., 1988a). Subsequently, we showed that similar increases in two of the three hippocampal mRNAs are elicited by the physiological stress of shaking for 2 h in a cage mounted on a metabolic shaker (Nichols et al., 1989). More recently, we have extended these studies to the cloned cDNA CORT-responses: GPDH/35 kd as an oligodendrocyte marker; CR16 as an unidentified gene expressed in hippocampal neurons; and proteolipid protein (PLP) as a non-changing control (Masters et al., 1987). Fig. 1 shows the results of RNA titration experiments on hippocampal total RNA prepared from individual rats in three treatment groups with the following serum CORT levels (ng/ml): ADX + stress, <10 ; intact, $<10 - 51$; intact + stress, 210 ± 40 . Although both GPDH and CR16 mRNAs increase in response to CORT in a dose-dependent manner and show type II glucocorticoid receptor specificity (unpublished data; Masters et al., 1988), they differ in their response to stress. In contrast to the significant 3-fold increase in GPDH mRNA abundance after 2 h vibratory stress, the CR16 transcript does not change between intact and intact-stressed rats. However, CR16 unlike GPDH mRNA does change significantly between ADX (not shown) or ADX-stressed and intact rats with low A.M. levels of CORT.

In RNA slot blot hybridization studies, GFAP and GS mRNAs also did not respond to 2 h vibratory stress (not shown). The data collected to date on CORT and stress responses are summarized in Table 2. Only GPDH/35 kd and 20 kd were responsive to both 2 h CORT treatment (1 mg and 10 mg, respectively) and 2 h vibratory stress (Nichols et al., 1989). The lack of response to stress of the other RNAs, despite elevated CORT levels, is consistent with their slower response to exogenous CORT (>2 h but <8 h). One exception is the RNA coding for the 33 kd polypeptide, which does respond to 2 h CORT treatment but not to 2 h vibratory stress.

Although GFAP/50 kd did not respond to physiological elevation of CORT following 2 h vibratory stress, we have recently demonstrated that the gene is under physiological regulation by CORT, since it is disinhibited after 7 days (but not 3 days) ADX (Fig. 2). This response is similar albeit in the opposite direction to the results observed for CR16 mRNA where the difference between A.M. intact levels and ADX occurs as early as 1 day post-ADX (unpublished data). As expected, GFAP mRNA decreases within 8 h after CORT replacement in ADX rats (Nichols et al., in press).

CONCLUSIONS

We have demonstrated by hybrid selection of rat brain total RNA with CORT-responsive cDNA clones and subsequent *in vitro* translation/2-d gel analysis that two of the clones hybridize to mRNAs coding for the 50 and 35 kd polypeptides. Partial DNA sequences of both clones and comparison to known genes in Genbank have identified them, and hence their translation products, as GFAP and GPDH, respectively (Table 1). The RNAs coding for the 33 and 20 kd polypeptides remain unidentified. A translation product was not resolved for CR16 on 2-d gels with a pI gradient of 5.0 to 7.0, which is consistent with its extremely basic isoelectric point, predicted from translated coding region DNA sequence (unpublished data).

We have found three subsets of hippocampal mRNA responses to CORT and vibratory stress which differ in temporal and level-dependent aspects of CORT regulation (Figs. 1 and 2 and Table 2). Adrenal medullary cells release catecholamines and substance P differentially in response to two different stressful circumstances (Vaupel et al., 1988). However, several different stressors increase preproenkephalin A mRNA in rat hypothalamus (Yoshikawa et al., 1985; Lightman and Young, 1987). Examining a variety of stressors, chosen to differ in terms of CORT levels attained and activation pathways, will determine if the data in Table 2 represent a general or unique pattern of expression of CORT-dependent responses in hippocampus.

Additional experiments on kinetics of response, circadian variation, and chronic stress are also warranted in order to interpret differential stress response data. For example, no difference between ADX and A.M. intact levels of GPDH mRNA (Figs. 1) is consistent with GPDH being regulated solely by type II glucocorticoid receptors as a molecular stress response (Nichols et al., 1988b). Since low A.M. CORT levels significantly alter GFAP and CR16 expression (Figs. 1 and 2), they may also be regulated by type I receptors and modulate stress responsiveness (Masters et al., 1989; Ratka et al., 1989). Mechanisms of adaptation to stress in brain are likely to involve responses that are both mediated by glucocorticoids and opposed by them (McEwen and Brinton, 1987). In this context, elevation of GFAP is an early astrocytic response to brain injury; a decrease

in GFAP mRNA and protein (O'Callaghan et al., in press) by CORT may be a homeostatic, adaptive response (Munck et al., 1984; Nichols et al., in press).

In summary, these studies demonstrate the feasibility of using sensitive probes to examine changes in gene expression in hippocampus in response to hormonal and environmental stress. By examining their temporal cascades and altered patterns of expression, we may identify mechanisms by which the brain coordinates acute and chronic adaptive responses to stress.

REFERENCES

1. Dallman, M.F., Akana, S.F., Cascio, C.S., Darlington, D.N., Jacobson, L., and Levin, N., Regulation of ACTH: variations on a theme of B, Recent Prog. Horm. Res., 43 (1987) 113-173.
2. Lee, J.J. and Costlow, N.A., A molecular titration assay to measure transcript prevalence levels, Methods in Enzymology, 152 (1987) 633-648.
3. Lightman, S.L. and Young, W.S., III, Changes in hypothalamic preproenkephalin A mRNA following stress and opiate withdrawal, Nature, 328 (1987) 643-645.
4. Mason, P.J. and Williams, J.G., Hybridization in the analysis of recombinant DNA, In B.D. Hames and S.J. Higgins (Eds.), Nucleic Acid Hybridization: A Practical Approach, IRL Press, Oxford, 1985, pp. 113-138.
5. Masters, J.N., Nichols, N.R. and Finch, C.E., Isolation of cDNA clones from rat hippocampus responsive to glucocorticoids, Neurosci. Abstr., 13 (1987) 1085.
6. Masters, J.N., Nichols, N.R. and Finch, C.E., Differences in response to vibratory stress of two glucocorticoid regulated genes in rat hippocampus, Neurosci. Abstr., 14 (1988) 18.
7. Masters, J.N., Nichols, N.R., Osterburg, H.H. and Finch, C.E., Corticosterone regulated genes in the hippocampus: regulation by the type I or type II glucocorticoid receptor, Neurosci. Abstr., 15 (1989) 1129.
8. McEwen, B.S. and Brinton, R.E., Neuroendocrine aspects of adaptation, In E.R.de Kloet, V.M. Wiegant and D. de Wied (Eds.), Progress in Brain Research, Vol. 72, Elsevier Science, Amsterdam, 1987, pp. 11-26.
9. McEwen, B.S., deKloet, E.R., and Rostene, W., Adrenal steroid receptors and actions in the nervous system, Physiol. Rev., 66 (1986) 1121-1188.
10. Munck, A., Guyre, P.M. and Holbrook, N.J., Physiological functions of glucocorticoids in stress and their relation to pharmacological actions, Endocr. Rev., 5 (1984) 25-44.

11. Nichols, N.R., Lerner, S.P., Masters, J.N., May, P.C., Millar, S.L. and Finch, C.E., Rapid corticosterone-induced changes in gene expression in rat hippocampus display type II glucocorticoid receptor specificity, Mol. Endocrinol., 2 (1988a) 282-290.
12. Nichols, N.R., Masters, J.N. and Finch, C.E., Rapid increase in glycerol phosphate dehydrogenase poly(A)+RNA as a corticoid-dependent stress response in the adult rat brain, J. Cell Biochem., supplement 12D (1988b) 320.
13. Nichols, N.R., Masters, J.N., May, P.C., de Vellis, J. and Finch, C.E., Corticosterone-induced responses in rat brain RNA are also evoked in hippocampus by acute vibratory stress, Neuroendocrinology, 49 (1989) 40-46.
14. Nichols, N.R., Osterburg, H.H., Masters, J.N., Millar, S.L. and Finch, C.E., Messenger RNA for glial fibrillary acidic protein is decreased in rat brain following acute and chronic corticosterone treatment, Mol. Brain Res., in press.
15. O'Callaghan, J.P., Brinton, R.E., and McEwen, B.S., Glucocorticoids regulate the concentration of glial fibrillary acidic protein throughout the brain, Brain Res., in press.
16. Ratka, A., Sutanto, W., Bloemers, M., and de Kloet, E.R., On the role of brain mineralocorticoids (type I) and glucocorticoid (type II) receptors in neuroendocrine regulation, Neuroendocrinology, 50 (1989) 117-123.
17. Swanson, L.W., Sawchenko, P.E. and Lind, R.W., Regulation of multiple peptides in CRF parvocellular neurosecretory neurons: implication for the stress response, In Hokfelt, T., Fuxe, K. and Pernow, B., (Eds.), Prog. in Brain Res., Vol. 68, Elsevier Science, Amsterdam, 1986, pp. 169-190.
18. Vaupel, R., Jarry, H., Schlomer, H. and Wuttke, W., Differential response of substance P-containing subtypes of adrenomedullary cells to different stressors, Endocrinology, 123 (1988) 2140-2145.
19. Yoshikawa, K., Hong, J. and Sabol, S.L., Electroconvulsive shock increases preproenkephalin messenger RNA abundance in rat hypothalamus, Proc. Natl. Acad. Sci., 82 (1985) 589-593.




Fig. 1. Hippocampal total RNA was isolated from individual intact non-stressed rats or intact and ADX rats which were given vibratory stress for 2 h in a cage mounted on a shaker. For RNA titration experiments, an excess of ^{32}P -labelled antisense RNA synthesized from subcloned cDNA inserts was hybridized to increasing concentrations of total RNA, followed by RNase A and T1 digestion and acid precipitation. Changes in the abundance of GPDH, CR16 and PLP RNAs in individual rats of each treatment group were calculated from the slopes of titration curves. The mean \pm SE are shown as histograms and significance levels (ANOVA) are versus non-stressed intact.




Fig. 2. Hippocampal total RNA was isolated from individual rats that were intact or ADX for 7 or 3 days and sacrificed in the morning. RNA samples (1 μ g/lane) were size fractionated on formaldehyde/agarose gels, transferred to nylon and hybridized with an antisense rat GFAP or PLP 32 P-cRNA probe. The total integrated density of a band was determined by computerized densitometry with automatic background subtraction. The mean \pm SE are shown as histograms and significance levels (ANOVA) are versus intact.

Table 1

IDENTIFICATION OF CORT-RESPONSIVE CLONES/TRANSLATION PRODUCTS

CLONE	GENE	TRANSLATION PRODUCT	
		MOL. WT.	pI
CR 1	GLUTAMINE SYNTHETASE	43 kd	6.7
CR 3	GLYCEROL PHOSPHATE DEHYDROGENASE	35 kd*	6.3
CR 16	UNKNOWN	NOT FOUND	
CR 46/59	GLIAL FIBRILLARY ACIDIC PROTEIN	50 kd*	5.5
CR 62	UNKNOWN	42 kd	5.4
	UNKNOWN	33 kd*	6.5
	UNKNOWN	20 kd*	6.9

*Previous CORT-responsive RNA translation products (Nichols et al., 1988)

Table 2

SUMMARY OF DIFFERENTIAL RESPONSES TO STRESS

RESPONSE	FOLD-CHANGE (TMT/CONTROL)	
	2 HR CORT	2 HR VIBRATORY STRESS
GPDH/35 kd	3	3
20 kd	10	3
33 kd	27	1
CR 16	1	1
GFAP/50 kd	1	1
GS/43 kd	1	1